Journal of Surgical Oncology

Table of Contents

Volume 87, Number 1, July 15, 2004

Guest Editorial: Telomerase: A Potential Molecular Marker and Therapeutic Target for Cancer
Joe Y. Chang 1

Variability in Axillary Lymph Node Dissection for Breast Cancer
Michael Schaapveld, Renée Otter, Elisabeth G.E. de Vries, Vaclav Fidler, Joris A.K. Grond, Winette T.A. van der Graaf, Pieter L. de Vogel, and Pax H.B. Willemse 4
The regional variation in the number of reported axillary lymph nodes and its effect on the pathological axillary nodal status for breast cancer were analyzed using population-based cancer registry data.

Clinical Relevance of Parasternal Uptake in Sentinel Node Procedure for Breast Cancer
Hans F.J. Fabry, Pim G.N.J. Mutsaers, Sybren Meijer, Hans Torrenga, Rik Pijpers, Paul A.M. Van Leeuwen, and Joost R.M. Van der Sijp 13
Patients with recurrent disease were reviewed. The status of the internal mammary nodes did not contribute to the decision for adjuvant treatment.

The Relationship Between Prognostic Factors of Breast Cancer and Thyroid Disorders in Turkish Women
Ömer Cengiz, Betül Bozkurt, Bülent Ünal, Osman Yildirim, Melih Karabeyoğlu, Abdullah Eroğlu, Belma Koçer, and Murat Ulaş 19
The incidence of thyroid disorders in breast cancer patients is investigated, along with the clinical and laboratory features of breast cancer and their relation to thyroid disorders.

Use of Autologous Instead of Allogeneic Blood Transfusion During Esophagectomy Prolongs Disease-Free Survival Among Patients With Recurrent Esophageal Cancer
Satoru Motoyama, Manabu Okuyama, Michihiko Kitamura, Reijiro Saito, Shuichi Kamata, Katsuyuki Murata, and Jun-ichi Ogawa 26
The present study was conducted with the aim of understanding better the clinical characteristics of recurrent esophageal cancer and determining whether any survival advantage is conferred by transfusing autologous instead of allogeneic blood during the esophagectomy. Autologous blood transfusion prolongs disease-free survival of esophageal cancer patients with recurrent malignancy following esophagectomy.

Detection of Sentinel Lymph Nodes in Gastric Cancers Based on Immunohistochemical Analysis of Micrometastases
Tomoaki Karube, Takenori Ochiai, Hideaki Shimada, Takashi Nikaidou, and Hideki Hayashi 32
Immunohistochemical analysis of micrometastases in sentinel and non-sentinel nodes of gastric cancer was performed. A high accuracy rate indicated the feasibility of SN biopsy as an indicator for less invasive treatment of gastric cancer.

Surgical Management and Outcome of Metachronous Krukenberg Tumors From Gastric Cancer
Jae Ho Cheong, Woo Jin Hyung, Jian Chen, Junuk Kim, Seung Ho Choi, and Sung Hoon Noh 39
The present study suggests that a metastasectomy may significantly improve the overall and progression free survival of patients with a metachronous Krukenberg tumor after gastric cancer provided that no gross residual disease remains. To define the patient group who benefits most from resection, the extent of disease and resectability must be carefully evaluated before surgery.

Adjuvant, Adoptive Immunotherapy With Tumor Infiltrating Lymphocytes Plus Interleukin-2 After Radical Hepatic Resection for Colorectal Liver Metastases: 5-Year Analysis
Andrea Gardini, Giorgio Ercolani, Angela Riccobon, Matteo Ravaioli, Laura Ridolfi, Emanuela Flamini, Ruggero Ridolfi, Gian Luca Grazi, Antonino Cavallari, and Dino Amadori 46
We selected two groups of patients (treatment and control arms) to assess the clinical usefulness and biological features of adoptive immunotherapy with TIL plus IL-2 for colorectal liver metastases after radical hepatic surgery. No significant benefits in overall or disease-free survival were found between the two groups after 5 years' surveillance. The biological results indicated a significant restoration of signal transmission molecules in TIL after IL-2 exposure, particularly in disease-free patients.

Adenocarcinoma Arising in Association With Gastric Heterotopic Pancreas: A Case Report and Review of the Literature
Lyska Emerson, Lester J. Layfield, L. Ralph Rohr, and Merril T. Dayton 53

How I Do It: A Useful Technique of Paramedian Skin Incision for Midline Laparotomy in Patients Requiring Colostomy in Left Lower Abdomen
Pankaj Chaturvedi, Sajid S. Quereshi, and Bhavin Shah 58

Volume 87, Number 2, August 1, 2004

Jewish Ethnicity and Black Race: Contrasting Influences on the Prognosis of Breast Cancer
William L. Donegan and Judy A. Tjoe 61
For reasons that are incompletely understood, race and ethnicity are associated with differing mortality from breast cancer. This investigation was undertaken to evaluate racial and ethnic influences on survival of women with breast cancers and to identify influential factors.

Esophageal Cancer: Outcomes of Surgery, Neoadjuvant Chemotherapy, and Three-Dimension Conformal Radiotherapy
Éric Fréchette, David A. Buck, Brian J. Kaplan, Theodore D. Chung, James E. Shaw, Lisa A. Kachnic, and James P. Neifeld 68
A retrospective review was performed on 118 patients treated for localized esophageal cancer between 1995 and 2002 at a single institution. A 47% pathologic complete response was noted with neoadjuvant chemoradiotherapy and was associated with a statistically significant increased 3-year survival (59% vs. 20%).

IV-DSA Is a New Diagnostic Tool for Axillary Lymph Node Metastasis in Breast Cancer Patients
Osamu Watanabe, Tadao Shimizu, Hiroshi Imamura, Jun Kinoshita, Yoshihito Utada, Toshihiro Okabe, Kiyomi Kimura, Akira Hirano, Motohiko Aiba, and Kenji Ogawa 75
The aim of this study is to know whether IV-DSA is useful to detect axillary lymph node metastasis of breast cancer and to evaluate the anigiogenesis of lymph nodes in the axilla.

Intrahepatic Cholangiocarcinoma Diagnosed Preoperatively as Hepatocellular Carcinoma
Masakazu Yamamoto, Shunichi Ariizumi, Takehito Otsubo, Hideo Katsuragawa, Satoshi Katagiri, Masayuki Nakano, and Ken Takasaki 80
Patients with intrahepatic cholangiocarcinoma given a preoperative diagnosis of hepatocellular carcinoma had distinct clinical features and could be treated with the same operation as hepatocellular carcinoma patients.

Commentary 83

Extent of Hepatic Resection Does Not Correlate With Toxicity Following Adjuvant Chemotherapy
Waldemar F. Carlo, Amanda J. Hummer, Lawrence Schwartz, Deidre Sullivan, Mithat Gonen, William Jarnagin, Yuman Fong, and Nancy Kemeny 85
This article will address the question of whether large hepatic resections resulting in a greater loss of healthy liver predisposes patients to developing toxicity from the subsequent chemotherapeutic regimens.

Incidentally Discovered Gallbladder Cancer: Role of Cryotherapy
Stephanie R. Goldberg and James P. Neifeld 91
Cryotherapy to the gallbladder fossa may represent a safe and effective alternative to hepatic resection in the treatment of gallbladder cancer.

Vascular Endothelial Growth Factor in Esophageal Cancer
Axel Kleespies, Markus Guba, Karl-Walter Jauch, and Christiane J. Bruns 95
This article reviews the prognostic significance of VEGF expression, microvessel density (MVD), and lymphangiogenic factors in squamous cell carcinoma (SCC), Barrett's dysplasia, and adenocarcinoma (AC) of the esophagus, their predictive value for treatment response to chemo-radiotherapy and new anti-angiogenic treatment strategies.

Commentary on Barranger et al., Contralateral Axillary Lymph Node Drainage in Breast Cancer: A Case Report: Unusual Lymph Drainage Pathways in Patients With Breast Cancer
Omgo E. Nieweg, Maartje C. van Rijk, Pieter J. Tanis, and Susanne H. Estourgie 105

Letter to the Editor: Re: Gipponi M, Solari N, Di Somma FC, et al.: New Fields of Application of the Sentinel Lymph Node Biopsy in the Pathologic Staging of Solid Neoplasms: Review of Literature and Surgical Perspectives. J Surg Oncol 2004: 85:171–179
Bin K. Kroon, Simon Horenblas, and Omgo E. Nieweg 107

Volume 87, Number 3, September 1, 2004

Guest Editorial: The World Federation of Surgical Oncology Societies: The Global Mission
Niall O'Higgins 109

Intra-Abdominal Metastases From Soft Tissue Sarcoma
Kasim A. Behranwala, Pratik Roy, Victoria Giblin, Roger A'Hern, Cyril Fisher, and J. Meirion Thomas 116
The prognostic significance of abdominal metastasis of STS is evaluated.

Primary Sarcoma of the Breast
Manoj Pandey, Aleyamma Mathew, Elizabeth K. Abraham, and Balakrishnan Rajan 121
A retrospective analysis is reported of 19 cases over a 21-year period.

Clear Cell Sarcoma in the Era of Sentinel Lymph Node Mapping
Waddah B. Al-Refaie, Mir W. Ali, David Z. Chu, Isaac B. Paz, and Sarah L. Blair 126
The authors report their experience using SLNB as an additional tool in the staging of clear cell sarcomas.

Identification of a Single Nucleotide Polymorphism in the MMP-1 Promoter in Chondrosarcoma
Ian D. Dickey and Sean P. Scully 130
The presence and prevalence of an SNP is determined, with an attempt to correlate this with clinical outcome.

Clinical Patterns and Treatment Outcome of Elderly Patients in Clinical Stage IB/II Non-Small Cell Lung Cancer
Tatsuro Okamoto, Riichiroh Maruyama, Fumihiro Shoji, Jiro Ikeda, Tetsuya Miyamoto, Tomomi Nakamura, Hiroshi Asoh, and Yukito Ichinose 134
The present paper demonstrates initial treatment patterns and outcomes of elderly patients with NSCLC in clinical stage IB, IIA, IIB. Radiotherapy is an alternative strategy for patients who cannot undergo surgery, especially with adenocarcinoma.

Management of Hepatocellular Carcinoma in Renal Transplant Recipients
Kenneth S. Chok, Chi Ming Lam, Fu Keung Li, Kelvin K. Ng, Ronnie T. Poon, Chung Mau Lo, and Sheung Tat Fan 139
The management and outcome of HCC in 5 renal transplant recipients were reviewed. Regular surveillance of the renal transplant recipients in HBV endemic areas with close collaboration between the renal physicians and the liver surgeons is the key point for improved outcome.

Case Report: Anti-Ri Antibody Opsoclonus-Myoclonus Syndrome and Breast Cancer: A Case Report and a Review of the Literature
David A. Weizman and Wey L. Leong 143

Review Article: Kaposi's Sarcoma: An Update
Robert A. Schwartz 146

Volume 87, Number 4, September 15, 2004

Guest Editorial: Rationale for Superficial Injection Techniques in Lymphatic Mapping in Breast Cancer Patients
Omgo E. Nieweg, Susanne H. Estourgie, Maartje C. van Rijk, and Bin. B.R. Kroon 153

Survival of Patients With Untreated Rectal Cancer
A. Melissa Solum, Robert H. Riffenburgh, and Peter A.S. Johnstone 157
A review of archival literature documenting outcomes in patients with untreated rectal cancer revealed relatively long survival for patients is possible. This important finding can best serve as a comparison to which new treatments can be fully evaluated.

Utility of CEA and CA 19-9 Tumor Markers in Diagnosis and Prognostic Assessment of Mucinous Epithelial Cancers of the Appendix
C. Pablo Carmignani, Regina Hampton, Christina E. Sugarbaker, David Chang, and Paul H. Sugarbaker 162
Tumor markers CEA and CA 19-9 have selected uses in gastrointestinal cancer. Data shown in this article suggest that they have a practical use in patients with primary and recurrent peritoneal dissemination of an epithelial appendiceal malignancy.

Risk Factors for Peritoneal Dissemination of Colorectal Cancer
Shung-Haur Yang, Jen-Kou Lin, Chiung-Ru Lai, Chien-Chih Chen, A. Fen-Yau Li, Wen-Yih Liang, and Jeng-Kae Jiang 167
In decreasing order, CA19-9, pT4, and age were significant risk factors of peritoneal dissemination of colorectal cancer.

Heparanase Expression in Human Colorectal Cancer and Its Relationship to Tumor Angiogenesis, Hematogenous Metastasis, and Prognosis
Tomohiro Sato, Akio Yamaguchi, Takanori Goi, Yasuo Hirono, Kazuo Takeuchi, Kanji Katayama, and Shigeru Matsukawa 174
Heparanase expression may have an important role in invasion and hematogenous metastasis, and may be a biologic marker of prognostic significance in colorectal cancer patients. In addition, we indicated that heparanase expression has an axial role in the tumor angiogenesis.

Comparison of Methylation-Specific Polymerase Chain Reaction (MSP) With Reverse Transcriptase-Polymerase Chain Reaction (RT-PCR) in Peripheral Blood of Gastric Cancer Patients
Hiroshi Koike, Daisuke Ichikawa, Hisashi Ikoma, Eigo Otsuji, Kazuya Kitamura, and Hisakazu Yamagishi 182
A detection rate of methylation-specific polymerase chain reaction (MSP) was higher than that of reverse transcriptase-polymerase chain reaction (RT-PCR) in serum of patients with gastric cancer, especially in early-stage disease. Both assays can serve as markers that allow selection of those cases requiring more intensive screening and aggressive postoperative treatment.

Case Report: FAP With Concurrent Duodenal Adenomatous Polyposis and Carcinoid Tumor
E. Ramsay Camp, Steven N. Hochwald, and Chen Liu 187

Review Article: Unknown Primary
Mitchell I. Chorost, M. Catherine Lee, Cindy B. Yeoh, Manuel Molina, and Bimal C. Ghosh 191

Author Index to Volume 87 205

Subject Index to Volume 87 207